


AD A0 46582


MSNW



# KINETIC STUDIES FOR XeF AND KrF LASERS

Prepared for  
OFFICE OF NAVAL RESEARCH

September 1977



## DISTRIBUTION STATEMENT A

Approved for public release  
Distribution Unlimited

AD No. \_\_\_\_\_  
DDC FILE COPY



**MATHEMATICAL SCIENCES NORTHWEST, Inc.**

11  
14 Sep 1977

Program Code Number TE20

14 MSNW-1062

12 14p.

12

9  
#and#  
QUARTERLY RPT. STATUS REPORT, NO. 3

6  
TITLE: KINETIC STUDIES FOR XeF AND KrF LASERS.

15  
Contract Number: N00014-76-C-1066, DARPA Order-1806


Effective Date: 1 September 1976

Expiration Date: 30 November 1977

Amount: \$99,909

Sponsored by

Defense Advanced Research Projects Agency  
DARPA Order No. 1806



10  
Principal Investigator: R. E. /Center (206) 827-0460  
C. H. /Fisher (206) 827-0460

Scientific Officer: Director, Physics Programs  
Physical Sciences Division  
Office of Naval Research  
Department of the Navy  
800 North Quincy Street  
Arlington, Virginia 22117

MATHEMATICAL SCIENCES NORTHWEST, INC.  
P. O. Box 1887  
Bellevue, Washington 98009

The views and conclusions contained in this document are those of the authors and should not be interpreted as necessarily representing the official policies, either expressed or implied, of the Defense Advanced Research Projects Agency or the U. S. Government.

DISTRIBUTION STATEMENT A

Approved for public release;  
Distribution Unlimited

40.4 058

Amc

## TABLE OF CONTENTS

SECTION		PAGE
I	INTRODUCTION	1
II	EXPERIMENTAL RESULTS	2
	2.1. XeF(B $\frac{1}{2}$ ) Radiative Lifetime	2
	2.2. XeF(B $\frac{1}{2}$ ) Collisional Quenching	4
	REFERENCES	9



# LIST OF FIGURES

FIGURE		PAGE
1	The 353 nm Fluorescence from a Gas Mixture of He + 10 Torr Xe + 0.7 Torr UF <sub>6</sub> at a Total Pressure of 125 Torr	3
2	Decay Frequency as a Function of Pressure for a He + 4% Xe + 0.3% UF <sub>6</sub> Gas Mixture	5
3	Variation of 353 nm Fluorescence Decay as a Function of Fluorine Partial Pressure	6

ACCESSION #		
NTIS	White Section	<input checked="" type="checkbox"/>
ODC	Defi Section	<input type="checkbox"/>
UNANNOUNCED		<input type="checkbox"/>
JUSTIFICATION		
<i>Letter on file</i>		
BY		
DISTRIBUTION/AVAILABILITY CODES		
Dist.	AVAIL.	and/or SPECIAL
<i>A</i>		

# LIST OF TABLES

TABLE

PAGE

1

Summary of  $\text{XeF}(B^{1/2})$  Radiative Lifetime and  
Quenching Data

7

## I. INTRODUCTION

The goal of this program is to determine the radiative lifetime and the collisional quenching rate coefficients for the XeF and KrF upper laser levels using a state selective, laser excited fluorescence technique. Details of the experimental procedure were described in our semi-annual report.<sup>†</sup> During this reporting period, we have determined the XeF( $B^{1/2}$ ) radiative lifetime by monitoring the decay of the 353 nm XeF( $B^{1/2} \rightarrow X$ ) fluorescence as a function of pressure. Preliminary collisional quenching rate constants have been determined for He, Xe, and F<sub>2</sub> from the variation of the fluorescence decay time constant as a function of partial pressure of the collision partner. Collisional quenching by Ar was investigated; however, no bimolecular rate constant was determined because of nonexponential decay of the XeF( $B^{1/2} \rightarrow X$ ) emission in the presence of Ar. This behavior suggests collisional coupling to some other excited state, most likely the XeF( $C^{3/2}$ ) state.

Bimolecular collisional quenching rate coefficients by Ne and NF<sub>3</sub> remain to be determined during the next reporting period. In addition, termolecular quenching by He, Ne, and Ar will be investigated by increasing the partial pressures of these constituents to the 1 to 3 atm range. Collisional coupling between the  $B^{1/2}$  and  $C^{3/2}$  states will also be investigated by looking for fluorescence at 460 nm on the XeF( $C \rightarrow X$ ) band.

## II. EXPERIMENTAL RESULTS

### 2.1. XeF(B $\frac{1}{2}$ ) Radiative Lifetime

Oscilloscope traces showing the 351 nm exciting pulse and the 353 nm fluorescence signal are presented in Figure 1. This signal was obtained with a He + 10 torr Xe + 0.7 torr UF<sub>6</sub> gas mixture at 125 torr total pressure. We have switched to UF<sub>6</sub> as the F atom source because its UV absorption cross section (resulting in dissociation to produce an F atom) is two orders of magnitude larger than that of F<sub>2</sub>, which yields a substantial increase in the XeF fluorescence. However, the UF<sub>6</sub> produced during the flash has a low vapor pressure resulting in the formation of solid particles which increased the scattered light from the 351 nm exciting pulse on subsequent shots. Therefore, it was necessary to pump out the cell and refill it with fresh gas after every shot.


If the XeF(B  $\frac{1}{2}$ ) excited state is lost only by radiative decay and bimolecular collisions, it will decay exponentially once the exciting laser pulse has terminated. A semilogarithmic plot of the 353 nm fluorescence decay read from the oscilloscope trace is presented in Figure 1. The slope of the straight line fit to the data yields the decay time constant  $\tau^{-1}$  given by

$$\begin{aligned}\tau^{-1} &= \tau_B^{-1} + k_{\text{He}} [\text{He}] + k_{\text{Xe}} [\text{Xe}] + k_{\text{UF}_6} [\text{UF}_6] \\ &= \tau_B^{-1} + (k_{\text{He}} X_{\text{He}} + k_{\text{Xe}} X_{\text{Xe}} + k_{\text{UF}_6} X_{\text{UF}_6}) p\end{aligned}$$


where  $\tau_B$  is the radiative lifetime of the XeF(B  $\frac{1}{2}$ ) state,  $k_i$  and  $X_i$  are the quenching rate coefficient and the mole fraction of the  $i^{\text{th}}$  component of the mixture, and  $p$  is the total pressure. Thus, a plot of the decay time constant  $\tau^{-1}$  as a function of pressure for a



a)



b)



77 01035

Figure 1. The 353 nm Fluorescence from a Gas Mixture of He+10 Torr Xe + 0.7 torr  $\text{UF}_6$  at a Total Pressure of 125 torr. a) Oscilloscope traces showing the 351 nm exciting laser pulse and the 353 nm fluorescence signal. b) Semi-logarithmic plot of the 353 nm fluorescence decay.




gas mixture with fixed composition should follow a straight line with the  $\text{XeF}(\text{B } \frac{1}{2})$  radiative lifetime as the zero pressure intercept. A plot of the data obtained for a He + 4% Xe + 0.13%  $\text{UF}_6$  gas mixture is shown in Figure 2. The zero pressure intercept of the straight line fit to the data gives a  $\text{XeF}(\text{B } \frac{1}{2})$  radiative lifetime of  $18.7 \pm 1.4$  nsec. This result is in good agreement with that obtained by Burnham and Harris from the dissociative excitation of  $\text{XeF}_2$ .<sup>2</sup>

## 2.2. $\text{XeF}(\text{B } \frac{1}{2})$ Collisional Quenching


Bimolecular quenching rate constants for various molecules are obtained by measuring the  $\text{XeF}(\text{B } \frac{1}{2})$  decay time constant as a function of the partial pressure of that constituent while holding the partial pressures of the other constituents fixed. The slope of the straight line fit to the data gives the two-body quenching rate constant for that molecule. To illustrate the procedure, a plot of the 353 nm decay time constant  $\tau^{-1}$  as a function of  $\text{F}_2$  partial pressure is shown in Figure 3. The partial pressures of the other constituents of the gas mixtures were held fixed at 250 torr He + 10 torr Xe + 0.3 torr  $\text{UF}_6$ . The quenching rate constant for  $\text{F}_2$  determined from the slope of the straight line fit to the data is  $k_{\text{F}_2} = 1.2 \times 10^{-10} \text{ cm}^3 \text{ molecule}^{-1} \text{ sec}^{-1}$ . Using this same procedure, quenching rate constants for He and Xe have also been determined. These rate constants are listed in Table 1 along with the pressure range over which the quenching was studied. The measured rate constants are in fairly good agreement with those inferred by Brashears, Setser and Desmarteau from Stern-Volmer plots of the quenching of  $\text{XeF}(\text{B } \frac{1}{2} \rightarrow \text{X})$  emission produced by photolysis of  $\text{XeF}_2$ .<sup>3</sup>

No quenching rate constant is listed in the table for Ar even though it is a primary constituent of e-beam excited XeF laser



77 01036

Figure 2. Decay Frequency as a Function of Pressure for a He + 4% Xe + 0.13% UF<sub>6</sub> Gas Mixture. The zero pressure intercept of the straight line fit to the data gives a XeF (B<sub>1/2</sub>) radiative lifetime of 18.7 nsec.



77 01037

Figure 3. Variation of 353 nm Fluorescence Decay as a Function of Fluorine Partial Pressure. The slope of the straight line fit to the data gives a quenching rate constant of  $1.2 \times 10^{-10} \text{ cm}^3 \text{ molecule}^{-1} \text{ sec}^{-1}$ .



Table 1

Summary of XeF (B  $\frac{1}{2}$ ) Radiative Lifetime  
and Quenching Data

XeF(B  $\frac{1}{2}$ )  $\tau_r = 18.7 \pm 1.4$  nsec  
XeF(B  $\frac{1}{2}$ ) 2-Body Quenching Rate Constants

Molecule	$k_q(\text{cm}^3 \text{ Molecule}^{-1} \text{ sec}^{-1})$	Pressure Range (torr)
He	$2 \times 10^{-12}$	75 - 750
Xe	$6 \times 10^{-11}$	5 - 80
F <sub>2</sub>	$1.2 \times 10^{-10}$	0.5 - 24

mixtures. This is due to the fact that the 353 nm fluorescence in the presence of Ar exhibited nonexponential decay over much of the pressure range studied, making it difficult to extract a rate constant without a more complicated analysis. This behavior suggests collisional transfer from the  $B^{1/2}$  state during the decay period. The most likely candidate is the  $XeF(C^{3/2})$  state which is nearly degenerate with the  $B^{1/2}$  state.<sup>4</sup> During the next reporting period, the possibility of collisional coupling between the  $B^{1/2}$  and  $C^{3/2}$  states will be investigated by looking for broadband fluorescence at 460 nm on the  $XeF(C^{3/2} \rightarrow X)$  transition.

#### REFERENCES

1. R.E. Center and C.H. Fisher, "Kinetic Studies for XeF and KrF Lasers," MSNW Semi-Annual Report, May 1977.
2. R. Burnham and N.W. Harris, J. Chem. Phys. 66, 2742 (1977).
3. H.C. Brashears, Jr., D.W. Setser, and D. Desmarteau, Chem. Phys. Lett. 48, 84 (1977).
4. T.H. Dunning and P.J. Hay, to be published.